

AN ANALOGUE OF ROW REMOVAL FOR DIAGRAMMATIC CHEREDNIK ALGEBRAS

C. BOWMAN AND L. SPEYER

ABSTRACT. We prove an analogue of James–Donkin row removal theorems for arbitrary diagrammatic Cherednik algebras. This is one of the first results concerning the (graded) decomposition numbers of these algebras over fields of arbitrary characteristic. As a special case, our results yield a new reduction theorem for graded decomposition numbers and extension groups for cyclotomic q -Schur algebras.

INTRODUCTION

Weighted Khovanov–Lauda–Rouquier algebras first arose in Webster’s work on Khovanov homology, knot theory, and higher representation theory [Web1]. Their finite dimensional quotients, the diagrammatic Cherednik algebras, have proven to be of great importance in the study of complex reflection groups, $G(\ell, 1, n)$, and their deformations [Webc].

Given an e -multicharge $\kappa \in (\mathbb{Z}/e\mathbb{Z})^\ell$ and a weighting $\theta \in \mathbb{R}^\ell$, we have an associated diagrammatic Cherednik algebra $A(n, \theta, \kappa)$. Over the complex field, the graded decomposition numbers of $A(n, \theta, \kappa)$ are now well-understood in terms of Uglov’s canonical bases of higher level Fock spaces [Los, RSVV, Webc]. In this paper and [BS], the authors initiate the study of the representation theory of these algebras over fields of arbitrary characteristic.

Given a weighting, $\theta \in \mathbb{R}^\ell$, Webster introduced a graphical calculus for the set of ℓ -multipartitions based on an embedding of these multipartitions into $\mathbb{R}^2$. We define **diagonal cuts** on these multipartitions in a graphical fashion; these cuts allow us to identify a pair of multipartitions (λ, μ) with two distinct pairs of multipartitions (λ^L, μ^L) and (λ^R, μ^R) (the left and right pieces of the cut) of smaller degree. We thus reduce the problem of calculating the graded decomposition numbers of these algebras as follows.

Main Theorem. *Let (λ, μ) be a pair of ℓ -multipartitions of n and let $a \in \mathbb{R}$. If (λ, μ) admits a θ -diagonal cut at $x = a$ into two pieces (λ^L, μ^L) and (λ^R, μ^R) , then we can factorise the graded decomposition numbers for these algebras as*

$$d_{\lambda\mu}(t) = d_{\lambda^L\mu^L}(t) \times d_{\lambda^R\mu^R}(t)$$

and the (graded) higher extension groups as

$$\mathrm{Ext}_{A(n, \theta, \kappa)}^k(\Delta(\lambda), \Delta(\mu)) \cong \bigoplus_{i+j=k} \mathrm{Ext}_{A(n_L, \theta, \kappa)}^i(\Delta(\lambda^L), \Delta(\mu^L)) \otimes \mathrm{Ext}_{A(n_R, \theta, \kappa)}^j(\Delta(\lambda^R), \Delta(\mu^R)),$$

where $n_L = |\lambda^L| = |\mu^L|$ and $n_R = |\lambda^R| = |\mu^R|$.

In the level $\ell = 1$ case, the algebra $A(n, \theta, \kappa)$ is Morita equivalent to the classical q -Schur algebra. In this case, one can make a diagonal cut across a pair of partitions if and only if one can make a **vertical** (generalised column) cut, if and only if one can make a **horizontal** (generalised row) cut. These vertical and horizontal cuts across pairs of partitions have already been extensively studied; the analogue of the above result for (graded) decomposition numbers was treated in [Jam81, Don85, CMT02] and for Ext groups in [FL03, LM05, Don07].

In higher levels, the (graphically defined) dominance orders on multipartitions are more exotic; our diagonal cuts may pass through many components of the multipartitions at once, as illustrated shortly. Our diagonal cuts provide new information even in the case of the cyclotomic q -Schur algebras of Dipper, James and Mathas [DJM98] (which are Morita equivalent to the diagrammatic Cherednik algebras with well-separated weightings); see Example 2.22.

In order to clarify the above, let's consider an example. For $g = 1$ and $\theta = (0, 0.5)$, the bipartitions

$$\lambda = ((11, 9, 7, 3^2, 2, 1^3), (9, 4, 2, 1^4)) \quad \text{and} \quad \mu = ((10, 9, 8, 4, 3, 1^5), (8, 4, 2, 1^4))$$

admit a θ -diagonal cut at $x = 2.6$, in the sense of Definition 2.1. To see this, we draw the bipartitions with respect to this weighting as in Figure 1. Roughly speaking, the left- and right-hand sides of these bipartitions with respect to this cut are the ‘smallest’ bipartitions which contain both the boxes intersected by the line and all the boxes to the left (respectively right). For λ and μ as above, these cuts are depicted in Figure 2. In this case,

$$\begin{aligned} \lambda^L &= ((11, 9, 7, 3^2), (9, 4, 2)), & \lambda^R &= ((3^5, 2, 1^3), (1^7)), \\ \mu^L &= ((10, 9, 8, 4, 3), (8, 4, 2)), & \mu^R &= ((3^5, 1^5), (1^7)). \end{aligned}$$

Now, applying our main theorem to this example, we obtain

$$d_{\lambda\mu}(t) = d_{\lambda^L\mu^L}(t) \times d_{\lambda^R\mu^R}(t).$$

Now let $e = 5$ and $\kappa = (0, 2)$. By results from [BS] (and the above) we obtain

$$d_{\lambda\mu}(t) = (t^5 + t^3) \times t^2 = t^7 + t^5.$$




FIGURE 1. We depict the bipartition $\lambda = ((11, 9, 7, 3^2, 2, 1^3), (9, 4, 2, 1^4))$ on the left and the bipartition $\mu = ((10, 9, 8, 4, 3, 1^5), (8, 4, 2, 1^4))$ on the right, with $g = 1$ and $\theta = (0, 0.5)$.




FIGURE 2. With bipartitions λ and μ as in Figure 1 and a diagonal cut at $x = 2.6$, we arrive at pairs of bipartitions (λ^L, μ^L) and (λ^R, μ^R) as depicted above.

Acknowledgements. The authors would like to thank the Royal Commission for the Exhibition of 1851 and the Japan Society for the Promotion of Science for their financial support.

1. THE DIAGRAMMATIC CHEREDNIK ALGEBRA

In this section we define the diagrammatic Cherednik algebras and recall the combinatorics underlying their representation theory.

Definition 1.1. Suppose that A is a $\mathbb{Z}$ -graded $\mathbb{k}$ -algebra which is of finite rank over $\mathbb{k}$. We say that A is a **graded cellular algebra with a highest weight theory** if the following conditions hold.

The algebra is equipped with a **cell datum** $(\Lambda, \mathcal{T}, C, \deg)$, where $(\Lambda, \triangleright)$ is the **weight poset**. For each $\lambda, \mu \in \Lambda$ such that $\lambda \triangleright \mu$, we have a finite set, denoted $\mathcal{T}(\lambda, \mu)$, and we let $\mathcal{T}(\lambda) = \cup_{\mu} \mathcal{T}(\lambda, \mu)$. There exist maps

$$C : \coprod_{\lambda \in \Lambda} \mathcal{T}(\lambda) \times \mathcal{T}(\lambda) \rightarrow A \quad \text{and} \quad \deg : \coprod_{\lambda \in \Lambda} \mathcal{T}(\lambda) \rightarrow \mathbb{Z}$$

such that C is injective. We denote $C(S, T) = c_{S,T}^\lambda$ for $S, T \in \mathcal{T}(\lambda)$. We require that A satisfies properties (1)–(6), below.

- (1) Each element $c_{S,T}^\lambda$ is homogeneous of degree

$$\deg(c_{S,T}^\lambda) = \deg(S) + \deg(T),$$

for $\lambda \in \Lambda$ and $S, T \in \mathcal{T}(\lambda)$.

- (2) The set $\{c_{S,T}^\lambda \mid S, T \in \mathcal{T}(\lambda), \lambda \in \Lambda\}$ is a $\mathbb{k}$ -basis of A .
 (3) If $S, T \in \mathcal{T}(\lambda)$, for some $\lambda \in \Lambda$, and $a \in A$ then there exist scalars $r_{S,U}(a)$, which do not depend on T , such that

$$ac_{S,T}^\lambda = \sum_{U \in \mathcal{T}(\lambda)} r_{S,U}(a) c_{U,T}^\lambda \pmod{A^{\triangleright \lambda}},$$

where $A^{\triangleright \lambda}$ is the $\mathbb{k}$ -submodule of A spanned by

$$\{c_{Q,R}^\mu \mid \mu \triangleright \lambda \text{ and } Q, R \in \mathcal{T}(\mu)\}.$$

- (4) The $\mathbb{C}$ -linear map $*$: $A \rightarrow A$ determined by $(c_{S,T}^\lambda)^* = c_{T,S}^\lambda$, for all $\lambda \in \Lambda$ and all $S, T \in \mathcal{T}(\lambda)$, is an anti-isomorphism of A .
 (5) The identity 1_A of A has a decomposition $1_A = \sum_{\lambda \in \Lambda} 1_\lambda$ into pairwise orthogonal idempotents 1_λ .
 (6) For $S \in \mathcal{T}(\lambda, \mu)$, $T \in \mathcal{T}(\lambda, \nu)$, we have that $1_\mu c_{S,T}^\lambda 1_\nu = c_{S,T}^\lambda$. There exists a unique element $T^\lambda \in \mathcal{T}(\lambda, \lambda)$, and $c_{T^\lambda, T^\lambda}^\lambda = 1_\lambda$.

All results in this section follow from [HM10]. Suppose that A is a graded cellular algebra with a highest weight theory. Given any $\lambda \in \Lambda$, the **graded standard module** $\Delta(\lambda)$ is the graded left A -module with basis $\{c_S^\lambda \mid S \in \mathcal{T}(\lambda, \mu), \mu \in \Lambda\}$. The action of A on $\Delta(\lambda)$ is given by

$$ac_S^\lambda = \sum_{U \in \mathcal{T}(\lambda)} r_{S,U}(a) c_U^\lambda,$$

where the scalars $r_{S,U}(a)$ are the scalars appearing in condition (3) of Definition 1.1. Suppose that $\lambda \in \Lambda$. There is a bilinear form $\langle \cdot, \cdot \rangle_\lambda$ on $\Delta(\lambda)$ which is determined by

$$c_{U,S}^\lambda c_{T,V}^\lambda \equiv \langle c_S^\lambda, c_T^\lambda \rangle_\lambda c_{U,V}^\lambda \pmod{A^{\triangleright \lambda}},$$

for any $S, T, U, V \in \mathcal{T}(\lambda)$.

If M is a graded A -module and $k \in \mathbb{Z}$, define $M\langle k \rangle$ to be the same module with $(M\langle k \rangle)_i = M_{i-k}$ for all $i \in \mathbb{Z}$. We call this a **degree shift** by k . If M is a graded A -module and L is a graded simple module let $[M : L\langle k \rangle]$ be the multiplicity of $L\langle k \rangle$ as a graded composition factor of M , for $k \in \mathbb{Z}$.

Suppose that A is a graded cellular algebra with a highest weight theory. For every $\lambda \in \Lambda$, define $L(\lambda)$ to be the quotient of the corresponding standard module $\Delta(\lambda)$ by the radical of the bilinear form $\langle \cdot, \cdot \rangle_\lambda$. This module is simple, and every simple module is of the form $L(\lambda)\langle k \rangle$ for

some $k \in \mathbb{Z}$, $\lambda \in \Lambda$. Let t be an indeterminate over $\mathbb{Z}_{\geq 0}$. The **graded decomposition matrix** of A is the matrix $\mathbf{D}_A(t) = (d_{\lambda\mu}(t))$, where

$$d_{\lambda\mu}(t) = \sum_{k \in \mathbb{Z}} [\Delta(\lambda) : L(\mu)\langle k \rangle] t^k,$$

for $\lambda, \mu \in \Lambda$.

1.1. Combinatorics of loadings. Fix integers $\ell, n \in \mathbb{Z}_{\geq 0}$, $g \in \mathbb{R}_{>0}$ and $e \in \{3, 4, \dots\} \cup \{\infty\}$. We define a **weighting** $\theta = (\theta_1, \dots, \theta_\ell) \in \mathbb{R}^\ell$ to be any ℓ -tuple such that $\theta_i - \theta_j$ is not an integer multiple of g for $1 \leq i < j \leq \ell$. Let κ denote an e -**multicharge** $\kappa = (\kappa_1, \dots, \kappa_\ell) \in (\mathbb{Z}/e\mathbb{Z})^\ell$.

An ℓ -**multipartition** $\lambda = (\lambda^{(1)}, \dots, \lambda^{(\ell)})$ of n is an ℓ -tuple of partitions such that $|\lambda^{(1)}| + \dots + |\lambda^{(\ell)}| = n$. We will denote the set of ℓ -multipartitions of n by $\mathcal{P}_n^\ell$. Let $\lambda = (\lambda^{(1)}, \lambda^{(2)}, \dots, \lambda^{(\ell)}) \in \mathcal{P}_n^\ell$. The **Young diagram** $[\lambda]$ is defined to be the set

$$\{(r, c, m) \in \mathbb{N} \times \mathbb{N} \times \{1, \dots, \ell\} \mid c \leq \lambda_r^{(m)}\}.$$

We refer to elements of $[\lambda]$ as **nodes** (of $[\lambda]$ or λ). We define the **residue** of a node $(r, c, m) \in [\lambda]$ to be $\kappa_m + c - r \pmod{e}$.

We define the **Russian array** as follows. For each $1 \leq m \leq \ell$, we place a point on the real line at θ_m and consider the region bounded by half-lines at angles $3\pi/4$ and $\pi/4$. We tile the resulting quadrant with a lattice of squares, each with diagonal of length $2g$. For each node of $[\lambda]$ we draw a box in the Russian array. We place the first node, $(1, 1, m)$, of component m at θ_m on the real line, with rows going northwest from this node, and columns going northeast. The diagram is tilted ever-so-slightly in the clockwise direction so that the top vertex of the box (r, c, m) (that is, the box in the r th row and c th column of the m th component of $[\lambda]$) has x -coordinate $\theta_m + g(r - c) + (r + c)\epsilon$.

Remark 1.2. Here the tilt ϵ is chosen so that $n\epsilon$ is absolutely small with respect to

- the ghost distance g (so that $\epsilon \ll g/n$)
- the weighting (so that g does not divide any number in the interval $|\theta_i - \theta_j| + (-n\epsilon, +n\epsilon)$ for $1 \leq i < j \leq \ell$).

Therefore given any $x \in \mathbb{R}$, the interval $(x - \epsilon, x + \epsilon) \subset \mathbb{R}$ contains at most one node from the loading of a box of an ℓ -multipartition of n .

Remark 1.3. With the assumption of Remark 1.2 firmly in place, we will drop any mention of ϵ when speaking of $g \in \mathbb{R}_{>0}$, or the weighting, $\theta \in \mathbb{R}^\ell$.

We define a **loading**, $\mathbf{i}$, to be an element of $(\mathbb{R} \times (\mathbb{Z}/e\mathbb{Z}))^n$ such that no real number occurs with multiplicity greater than one. Given a node (r, c, m) in some $\lambda \in \mathcal{P}_n^\ell$, the associated **loading**, $\mathbf{i}_{(r, c, m)}$, is given by the pair

$$(\theta_m + g(r - c) + (r + c)\epsilon, \kappa_m + c - r) \in \mathbb{R} \times (\mathbb{Z}/e\mathbb{Z}),$$

where $\theta_m + g(r - c) + (r + c)\epsilon \in \mathbb{R}$ is the x -coordinate of the top vertex of the node (r, c, m) and $\kappa_m + c - r$ is the residue of the node (r, c, m) . Given $\lambda \in \mathcal{P}_n^\ell$, we define the **loading**, $\mathbf{i}_\lambda \in (\mathbb{R} \times (\mathbb{Z}/e\mathbb{Z}))^n$, to be the n -tuple which consists of the loadings of all n nodes in $[\lambda]$.

We let D_λ denote the underlying ordered subset of $\mathbb{R}$ given by the points of the loading of $\lambda \in \mathcal{P}_n^\ell$. Given $a \in D_\lambda$, we abuse notation and let a denote the corresponding node of λ (that is, the node whose top vertex projects onto x -coordinate $a \in \mathbb{R}$). The **residue sequence** of λ is given by reading the residues of the nodes of λ according to the ordering given by D_λ .

Example 1.4. Let $\ell = 2$, $g = 1$, $\epsilon = 1/100$, and $\theta = (0, 0.5)$. The bipartition $((2, 1), (1^3))$ has Young diagram and corresponding loading $\mathbf{i}_\lambda$ given in Figure 3. The residue sequence of λ is $(\kappa_1 + 1, \kappa_1, \kappa_2, \kappa_1 - 1, \kappa_2 - 1, \kappa_2 - 2)$, and

$$D_\lambda = \{-0.97, 0.02, 0.52, 1.03, 1.53, 2.54\}.$$

The node $x = -0.97$ in λ can be identified with the node in the first row and second column of the first component of λ .




FIGURE 3. The diagram and loading of the bipartition $((2, 1), (1^3))$ for $\ell = 2$, $g = 1$, $\theta = (0, 0.5)$.

1.2. Tableaux combinatorics. We now recall the tableaux combinatorics of diagrammatic Cherednik algebras, as defined in [Webc, Section 2.2].

Definition 1.5. Let $\lambda, \mu \in \mathcal{P}_n^\ell$. A λ -tableau of weight μ is a bijective map $T : [\lambda] \rightarrow D_\mu$ which respects residues. In other words, we fill a given node (r, c, m) of the diagram $[\lambda]$ with a real number d from D_μ (without multiplicities) so that the residue attached to the real number d in the loading $\mathbf{i}_\mu$ is equal to $\kappa_m + c - r \pmod{e}$.

Definition 1.6. A λ -tableau, T , of shape λ and weight μ is said to be **semistandard** if

- $T(1, 1, m) > \theta_m$,
- $T(r, c, m) > T(r - 1, c, m) + g$,
- $T(r, c, m) > T(r, c - 1, m) - g$.

We denote the set of all semistandard tableaux of shape λ and weight μ by $\text{SStd}(\lambda, \mu)$.

Definition 1.7. Let $\mathbf{i}$ and $\mathbf{j}$ denote two loadings of size n and let $r \in \mathbb{Z}/e\mathbb{Z}$. We say that $\mathbf{i}$ **r -dominates** $\mathbf{j}$ if for every real number $a \in \mathbb{R}$, we have that

$$|\{(x, r) \in \mathbf{i} \mid x < a\}| \geq |\{(x, r) \in \mathbf{j} \mid x < a\}|.$$

We say that $\mathbf{i}$ **dominates** $\mathbf{j}$ if $\mathbf{i}$ r -dominates $\mathbf{j}$ for every $r \in \mathbb{Z}/e\mathbb{Z}$. Given $\lambda, \mu \in \mathcal{P}_n^\ell$ and $\theta \in \mathbb{R}^\ell$, we say that λ **θ -dominates** μ (and write $\mu \leq_\theta \lambda$) if $\mathbf{i}_\lambda$ dominates $\mathbf{i}_\mu$.

1.3. The diagrammatic Cherednik algebra. Recall that we have fixed $\ell, n \in \mathbb{Z}_{>0}$, $g \in \mathbb{R}_{>0}$ and $e \in \{3, 4, \dots\} \cup \{\infty\}$. Given any weighting $\theta \in \mathbb{R}^\ell$ and any e -multicharge $\kappa \in (\mathbb{Z}/e\mathbb{Z})^\ell$, we now recall the definition of the diagrammatic Cherednik algebra, $A(n, \theta, \kappa)$.

This is an example of one of many finite dimensional algebras (‘reduced steadied quotients of weighted KLR algebras’ in Webster’s terminology) constructed in [Webc], whose module categories are equivalent, over the complex field, to category $\mathcal{O}$ for the rational cyclotomic Cherednik algebra [Webc, Theorem 3.9].

Definition 1.8. We define a θ -**diagram** of type $G(\ell, 1, n)$ to be a **frame** $\mathbb{R} \times [0, 1]$ with distinguished black points on the northern and southern boundaries given by the loadings $\mathbf{i}_\mu$ and $\mathbf{i}_\lambda$ for some $\lambda, \mu \in \mathcal{P}_n^\ell$ and a collection of curves each of which starts at a northern point and ends at a southern point of the same residue, *i* say (we refer to this as a **black i -strand**). We further require that each curve has a mapping diffeomorphically to $[0, 1]$ via the projection to the y -axis. Each curve is allowed to carry any number of dots. We draw

- a dashed line g units to the left of each strand, which we call a **ghost i -strand** or **i -ghost**;
- vertical red lines at $\theta_m \in \mathbb{R}$ each of which carries a residue κ_m for $1 \leq m \leq \ell$ which we call a **red κ_m -strand**.

We require that there are no triple points or tangencies involving any combination of strands, ghosts or red lines and no dots lie on crossings. We consider these diagrams equivalent if they are related by an isotopy that avoids these tangencies, double points and dots on crossings.

Definition 1.9 (Definition 4.1 [Webc]). The diagrammatic Cherednik algebra, $A(n, \theta, \kappa)$, is the $\mathbb{C}$ -algebra spanned by all θ -diagrams modulo the following local relations (here a local relation means one that can be applied on a small region of the diagram).

(2.1) Any diagram may be deformed isotopically; that is, by a continuous deformation of the diagram which at no point introduces or removes any crossings of strands (black, ghost, or red).

(2.2) For $i \neq j$ we have that dots pass through crossings.

$$\begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ i \quad j \end{array} = \begin{array}{c} \diagup \quad \diagdown \\ i \quad j \\ \bullet \end{array}$$

(2.3) For two like-labelled strands, we get an error term.

$$\begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ i \quad i \end{array} = \begin{array}{c} \diagup \quad \diagdown \\ i \quad i \\ \bullet \end{array} + \begin{array}{c} | \quad | \\ i \quad i \end{array} \quad \begin{array}{c} \diagup \quad \diagdown \\ i \quad i \\ \bullet \end{array} = \begin{array}{c} \diagup \quad \diagdown \\ i \quad i \\ \bullet \end{array} + \begin{array}{c} | \quad | \\ i \quad i \end{array}$$

(2.4) For double crossings of black strands, we have the following.

$$\begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ i \quad i \end{array} = 0 \quad \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ i \quad j \end{array} = \begin{array}{c} | \quad | \\ i \quad j \end{array}$$

(2.5) If $j \neq i - 1$, then we can pass ghosts through black strands.

$$\begin{array}{c} \text{ghost} \\ \diagup \quad \diagdown \\ i \quad j \end{array} \begin{array}{c} \diagup \quad \diagdown \\ i \quad j \end{array} = \begin{array}{c} | \quad | \\ i \quad j \end{array} \quad \begin{array}{c} \text{ghost} \\ \diagdown \quad \diagup \\ i \quad j \end{array} \begin{array}{c} \diagup \quad \diagdown \\ i \quad j \end{array} = \begin{array}{c} | \quad | \\ i \quad j \end{array}$$

(2.6) On the other hand, in the case where $j = i - 1$, we have the following.

$$\begin{array}{c} \text{ghost} \\ \diagup \quad \diagdown \\ i \quad i-1 \end{array} \begin{array}{c} \diagup \quad \diagdown \\ i-1 \quad i \end{array} = \begin{array}{c} | \quad | \\ i \quad i-1 \end{array} - \begin{array}{c} \bullet \\ | \\ i-1 \end{array} - \begin{array}{c} \bullet \\ | \\ i \end{array} \begin{array}{c} | \quad | \\ i-1 \quad i \end{array}$$

(2.7) We also have the relation below, obtained by symmetry.

$$\begin{array}{c} \text{ghost} \\ \diagdown \quad \diagup \\ i \quad i-1 \end{array} \begin{array}{c} \diagdown \quad \diagup \\ i-1 \quad i \end{array} = \begin{array}{c} | \quad | \\ i \quad i-1 \end{array} - \begin{array}{c} \bullet \\ | \\ i-1 \end{array} - \begin{array}{c} \bullet \\ | \\ i \end{array} \begin{array}{c} | \quad | \\ i-1 \quad i \end{array}$$

(2.8) Strands can move through crossings of black strands freely.

$$\begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ i \quad j \quad k \end{array} = \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ i \quad j \quad k \end{array}$$

Similarly, this holds for triple points involving ghosts, except for the following relations when $j = i - 1$.

(2.9)

$$\begin{array}{c} \text{ghost} \\ \diagup \quad \diagdown \\ i \quad j \end{array} \begin{array}{c} \diagup \quad \diagdown \\ j \quad j \end{array} = \begin{array}{c} \text{ghost} \\ \diagdown \quad \diagup \\ i \quad j \end{array} \begin{array}{c} \diagdown \quad \diagup \\ j \quad j \end{array} - \begin{array}{c} | \quad | \quad | \\ i \quad j \quad j \end{array}$$

(2.10)

$$\begin{array}{c} \text{ghost} \\ \diagdown \quad \diagup \\ i \quad i \end{array} \begin{array}{c} \diagdown \quad \diagup \\ j \quad j \end{array} = \begin{array}{c} \text{ghost} \\ \diagup \quad \diagdown \\ i \quad i \end{array} \begin{array}{c} \diagup \quad \diagdown \\ j \quad j \end{array} + \begin{array}{c} | \quad | \quad | \\ i \quad i \quad j \end{array}$$

The ghost strands may pass through red strands freely. For $i \neq j$, the black i -strands may pass through red j -strands freely. If the red and black strands have the same label, a dot is added to the black strand when straightening. Diagrammatically, these relations are given by

(2.11)

and their mirror images. All black crossings and dots can pass through red strands, with a correction term.

(2.12)

(2.13)

(2.14)

Finally, we have the following non-local idempotent relation.

(2.15) Any idempotent where the strands can be broken into two groups separated by a blank space of size $> g$ (so no ghost from the right-hand group can be left of a strand in the left group and *vice versa*) with all red strands in the right-hand group is referred to as unsteady and set to be equal to zero.

Remark 1.10. We remark that the θ -diagrams are clearly dependent on our choice of the parameters $e, \ell, n \in \mathbb{N}$ and $g \in \mathbb{R}_{>0}$, as well as $\theta \in \mathbb{R}^\ell$ and $\kappa \in (\mathbb{Z}/e\mathbb{Z})^\ell$. The dependence of $A(n, \theta, \kappa)$ on e and ℓ is subsumed by the dependence on κ . We have chosen to not emphasise the dependence of $A(n, \theta, \kappa)$ on $g \in \mathbb{R}_{>0}$ because this ‘ghost’ variable can always be scaled appropriately so that the definition of the algebra depends only on $\theta \in \mathbb{R}^\ell$. For a comparison of our notation with Webster’s, see [BCS, Section 2.2].

1.4. The grading on the diagrammatic Cherednik algebra. This algebra is graded as follows:

- dots have degree 2;
- the crossing of two strands has degree 0, unless they have the same label, in which case it has degree -2 ;
- the crossing of a black strand with label i and a ghost has degree 1 if the ghost has label $i - 1$ and 0 otherwise;
- the crossing of a black strand with a red strand has degree 0, unless they have the same label, in which case it has degree 1.

In other words,

$$\begin{aligned} \deg \bullet_i &= 2 & \deg \times_{i,j} &= -2\delta_{i,j} & \deg \times_{i,j}^{--} &= \delta_{j,i+1} & \deg \times_{i,j}^{--} &= \delta_{j,i-1} \\ \deg \times_{i,j}^{--} &= \delta_{i,j} & \deg \times_{i,j}^{--} &= \delta_{j,i}. \end{aligned}$$

1.5. A cellular basis of the diagrammatic Cherednik algebra. Let d be any θ -diagram and $y \in [0, 1]$ be any fixed value such that there are no crossings in d at any point in $\mathbb{R} \times \{y\}$. Then the positions of the various strands in this horizontal slice give a loading $\mathbf{i}_y$. We say that the diagram $d \in A(n, \theta, \kappa)$ factors through the loading $\mathbf{i}_y$.

The following lemma is a trivial consequence of the proof of [Webc, Lemma 2.15]. It is a key ingredient in the proof that $\mathcal{C}$ is a (cellular) basis of $A(n, \theta, \kappa)$ (see Theorem 1.12).

Lemma 1.11. *Suppose that a θ -diagram, d , factors through a loading $\mathbf{i}$ such that $\mathbf{i} \triangleright \mathbf{i}_\mu$ for some $\mu \in \mathcal{P}_n^\ell$, with $\mathbf{i}$ and $\mathbf{i}_\mu$ not isotopic. Then d factors through some $\mathbf{i}_\lambda$ with $\mathbf{i}_\lambda \triangleright \mathbf{i}$, for some $\lambda \in \mathcal{P}_n^\ell$.*

Given any $\mathbf{T} \in \text{SStd}(\lambda, \mu)$, we have a θ -diagram $B_{\mathbf{T}}$ consisting of a frame in which the n black strands each connecting a northern and southern distinguished point are drawn so that they trace out the bijection determined by $\mathbf{T}$ in such a way that we use the minimal number of crossings without creating any bigons between pairs of strands or strands and ghosts. This diagram is not unique up to isotopy (since we have not specified how to resolve triple points), but we can choose one such diagram arbitrarily.

Given a pair of semistandard tableaux of the same shape $(\mathbf{S}, \mathbf{T}) \in \text{SStd}(\lambda, \mu) \times \text{SStd}(\lambda, \nu)$, we have a diagram $C_{\mathbf{S}, \mathbf{T}} = B_{\mathbf{S}} B_{\mathbf{T}}^*$ where $B_{\mathbf{T}}^*$ is the diagram obtained from $B_{\mathbf{T}}$ by flipping it through the horizontal axis. Notice that there is a unique element $\mathbf{T}^\lambda \in \text{SStd}(\lambda, \lambda)$ and the corresponding basis element $1_\lambda = C_{\mathbf{T}^\lambda, \mathbf{T}^\lambda}$ is the idempotent in which all black strands are vertical. A degree function on tableaux is defined in [Webc, Definition 2.13]; for our purposes it is enough to note that $\deg(\mathbf{T}) = \deg(B_{\mathbf{T}})$ as we shall always work with the θ -diagrams directly.

The following theorem (see [Webc, Sections 2.6 and 4.4]) provides a cellular basis, $\mathcal{C}$, of the algebra $A(n, \theta, \kappa)$; the essential idea behind the proof is to rewrite any diagram $d \in A(n, \theta, \kappa)$ using relations (2.1) to (2.15) to push all black strands (and their ghosts) as far left as possible. Webster shows that this process of moving the strands leftwards eventually terminates with a linear combination of diagrams belonging to $\mathcal{C}$.

Theorem 1.12. *The algebra $A(n, \theta, \kappa)$ is a graded cellular algebra with a highest weight theory. The cellular basis is given by*

$$\mathcal{C} = \{C_{\mathbf{S}, \mathbf{T}} \mid \mathbf{S} \in \text{SStd}(\lambda, \mu), \mathbf{T} \in \text{SStd}(\lambda, \nu), \lambda, \mu, \nu \in \mathcal{P}_n^\ell\}$$

with respect to the θ -dominance order on the set $\mathcal{P}_n^\ell$ and the anti-isomorphism given by flipping a diagram through the horizontal axis.

Remark 1.13. We say that $\theta \in \mathbb{R}^\ell$ is a **well-separated weighting** for $A(n, \theta, \kappa)$ if $|\theta_j - \theta_i| > ng$ for all $1 \leq i < j \leq \ell$. In [Webb, Theorem 3.9] it is shown that if $\theta \in \mathbb{R}^\ell$ is well-separated, then the algebra $A(n, \theta, \kappa)$ is Morita equivalent to the corresponding cyclotomic q -Schur algebra.

We say that a weighting $\theta \in \mathbb{R}^\ell$ is a **FLOTW weighting** for $A(n, \theta, \kappa)$ if $0 < |\theta_i - \theta_j| < g$ for all $1 \leq i < j \leq \ell$.

2. DIAGONAL CUTS OF MULTIPARTITIONS

In this section, we identify subquotients of the diagrammatic Cherednik algebras which are isomorphic to the product of subquotients of two smaller diagrammatic Cherednik algebras. We first do this on the level of the indexing sets of representations (in Subsections 2.1 and 2.2) and then lift this to the level of algebras (Subsection 2.3). We show that the graded decomposition numbers and higher extension groups are preserved under taking these subquotients and hence prove the main theorem stated in the introduction.

2.1. Diagonal cuts and subquotients of the set of multipartitions. Let $a \in \mathbb{R}$ be such that the interval $(a - n\epsilon, a + n\epsilon) \subset \mathbb{R}$ does not contain the x -coordinate of the loading of a node belonging to any ℓ -multipartition of n . Given $\theta \in \mathbb{R}^\ell$ a weighting and $\lambda \in \mathcal{P}_n^\ell$, we define the set

$$I_a(\lambda) = \mathbf{i}_\lambda \cap ((a - g, a) \times \mathbb{Z}/e\mathbb{Z}).$$

That is, $I_a(\lambda)$ is the loading corresponding to all nodes $(r, c, m) \in [\lambda]$ which are slightly to the left of the line $x = a$; in other words,

$$(2.1) \quad \theta_m + g(r - c) < a < \theta_m + g(r - c) + g.$$

Note that for each $1 \leq m \leq \ell$, the set $I_a(\lambda) \cap \mathbf{i}_{\lambda^{(m)}}$ corresponds to a single diagonal in $[\lambda^{(m)}]$. Similarly, we define

$$L_a(\lambda) = \mathbf{i}_\lambda \cap ((-\infty, a - g) \times \mathbb{Z}/e\mathbb{Z}) \quad \text{and} \quad R_a(\lambda) = \mathbf{i}_\lambda \cap ((a, \infty) \times \mathbb{Z}/e\mathbb{Z}).$$

That is, $L_a(\lambda)$ (respectively $R_a(\lambda)$) is the loading corresponding to all nodes in $[\lambda]$ which are strictly to the left (respectively right) of the line $x = a - g$ (respectively $x = a$).

Definition 2.1. Let θ be a weighting, $\lambda, \mu \in \mathcal{P}_n^\ell$ and $a \in \mathbb{R}$. We say that the pair (λ, μ) admits a θ -diagonal cut at $x = a$ if the nodes in the region $(a - g, a)$ are common to both λ and μ , and the number of nodes strictly to the left or right of the line $x = a$ is the same in both multipartitions; in other words,

$$I_a(\lambda) = I_a(\mu), \quad |L_a(\lambda)| = |L_a(\mu)|, \quad \text{and} \quad |R_a(\lambda)| = |R_a(\mu)|.$$

Given $\lambda \in \mathcal{P}_n^\ell$, we let

$$\Lambda_a(n, \theta, \kappa) = \{\mu \in \mathcal{P}_n^\ell \mid (\lambda, \mu) \text{ admits a } \theta\text{-diagonal cut at } x = a\}.$$

Example 2.2. Let $e = 3$, $g = 2$, $\theta = 0$ and $(\lambda, \mu) = ((5, 4, 3, 2, 1), (4^3, 1^3))$. This pair admits a diagonal cut at $x = 1$. This cut is slightly to the right of the top vertex of the box $(3, 3)$. In this case

$$I_a(\lambda) = I_a(\mu) = \{((0 + 2\epsilon), 0), ((0 + 4\epsilon), 0), ((0 + 6\epsilon), 0)\}$$

and

$$|L_a(\lambda)| = |L_a(\mu)| = 6, \quad |R_a(\lambda)| = |R_a(\mu)| = 6.$$

Note that the pair (λ, μ) also admits a horizontal cut after the third row, and a vertical cut after the third column (in the sense of [Don85, FL03]). That is,

$$5 + 4 + 3 = \lambda_1 + \lambda_2 + \lambda_3 = \mu_1 + \mu_2 + \mu_3 = 4 + 4 + 4,$$

and

$$5 + 4 + 3 = \lambda'_1 + \lambda'_2 + \lambda'_3 = \mu'_1 + \mu'_2 + \mu'_3 = 6 + 3 + 3.$$




FIGURE 4. Fix $e = 3$, $g = 2$ and $\theta = 0$. The partitions $(5, 4, 3, 2, 1)$ and $(4^3, 1^3)$ admit a diagonal cut at $x = 1$.

Remark 2.3. Let (λ, μ) be a pair of partitions such that $\lambda \triangleright \mu$. The above example illustrates that in level 1, the pair (λ, μ) admits a θ -diagonal cut if and only if they admit a horizontal cut, if and only if they admit a vertical cut (in the sense of [Don85, FL03]).

Lemma 2.4. The set $\Lambda_a(n, \theta, \kappa)$ is closed under the θ -dominance order. That is, if $\mu, \mu' \in \Lambda_a(n, \theta, \kappa)$ and $\nu \in \mathcal{P}_n^\ell$ are such that $\mu \triangleright_\theta \nu \triangleright_\theta \mu'$, then $\nu \in \Lambda_a(n, \theta, \kappa)$.

Proof. By the definition of the θ -dominance order, the number of nodes to the left of the point $x = a - g$ for each of the three multipartitions is bounded as follows,

$$(2.2) \quad |L_a(\mu)| \geq |L_a(\nu)| \geq |L_a(\mu')|.$$

Recall (by the definition of $\Lambda_a(n, \theta, \kappa)$) that $|L_a(\mu')| = |L_a(\lambda)| = |L_a(\mu)|$; therefore $|L_a(\lambda)| = |L_a(\nu)|$, as required. Similarly, we have that the number of nodes to the right of $x = a$ for each of the multipartitions is bounded as follows,

$$(2.3) \quad |R_a(\mu')| \geq |R_a(\nu)| \geq |R_a(\mu)|.$$

Therefore (as above) we have that $|R_a(\nu)| = |R_a(\lambda)|$, as required.

It is immediate from equations (2.2) and (2.3), that $|I_a(\nu)| = |I_a(\lambda)|$. It remains to show that $I_a(\nu) = I_a(\lambda)$. To show this, we argue similarly to above, one node at a time, in the interval

$(a - g, a)$. First, we recall that our tilt ϵ is chosen to be small enough (with respect to θ and g ; see Remark 1.2) so that any interval $(x - \epsilon, x + \epsilon)$ contains the x -coordinate of at most one node of an element of $\mathcal{P}_n^\ell$. Therefore, if $x \in D_\lambda$ for some $\lambda \in \mathcal{P}_n^\ell$, there is precisely one such point in the interval $(x - \epsilon, x + \epsilon)$, namely x itself.

We note that if $D_\lambda \cap (a - g, a)$ is empty, then the result follows immediately from equations (2.2) and (2.3). Now, suppose that $D_\lambda \cap (a - g, a)$ is non-empty and take $x \in D_\lambda \cap (a - g, a)$. By the above arguments, we have that

$$|L_{x-\epsilon}(\mu)| = |L_{x-\epsilon}(\nu)| = |L_{x-\epsilon}(\mu')|.$$

Arguing as before, we must also have that

$$|L_{x+\epsilon}(\mu)| \geq |L_{x+\epsilon}(\nu)| \geq |L_{x+\epsilon}(\mu')|$$

where

$$|L_{x+\epsilon}(\mu)| = |L(\nu)_{x-\epsilon}| + 1 = |L_{x+\epsilon}(\mu')|.$$

Therefore, $L(\nu)$ has precisely one node in the open interval $(x - \epsilon, x + \epsilon)$. But the point x is the only node in this interval. So $x \in I_a(\nu)$, and the proof is complete. $\square$

Definition 2.5. Let P denote a partially ordered set and Q a subset of P . We say that Q is **saturated** if for any $\alpha \in Q$ and $\beta \in P$ with $\beta \triangleleft \alpha$, we have that $\beta \in Q$. We say that Q is **cosaturated** if its complement in P is saturated.

Given a subset $R \subseteq P$, we let $E(R)$ (respectively $F(R)$) denote the saturated (respectively cosaturated) closure of R in P ; that is, the smallest saturated (respectively cosaturated) set containing R .

By Lemma 2.4, we immediately deduce the following corollary.

Corollary 2.6. *We have that $\Lambda_a(n, \theta, \kappa) = E(\Lambda_a(n, \theta, \kappa)) \cap F(\Lambda_a(n, \theta, \kappa))$.*

Remark 2.7. In what follows, when the choice of $\lambda \in \mathcal{P}_n^\ell$, $a \in \mathbb{R}$, and $\theta \in \mathbb{R}^\ell$ is clear, we shall let $E := E(\Lambda_a(n, \theta, \kappa))$ and $F := F(\Lambda_a(n, \theta, \kappa))$.

2.2. A bijection between partially ordered sets. We now construct the bijection between partially ordered sets which will underly the isomorphism of subquotient algebras in Subsection 2.3.

Definition 2.8. Given $\lambda \in \mathcal{P}_n^\ell$, we define the left-hand side of λ (with respect to a θ -diagonal cut at $x = a$), denoted λ^L , to be the smallest ℓ -multipartition such that

$$I_a(\lambda) \cup L_a(\lambda) \subseteq \mathbf{i}_{\lambda^L}.$$

Similarly, we define the right-hand side of λ , denoted λ^R , to be the smallest ℓ -multipartition such that

$$I_a(\lambda) \cup R_a(\lambda) \subseteq \mathbf{i}_{\lambda^R}.$$

We let $n_L = |\lambda^L|$ and $n_R = |\lambda^R|$. We extend the notation of Definition 2.1 as follows,

$$\Lambda_a^L(n, \theta, \kappa) = \{\mu \in \mathcal{P}_{n_L}^\ell \mid (\lambda^L, \mu) \text{ admit a } \theta\text{-diagonal cut at } x = a\}$$

and

$$\Lambda_a^R(n, \theta, \kappa) = \{\mu \in \mathcal{P}_{n_R}^\ell \mid (\lambda^R, \mu) \text{ admit a } \theta\text{-diagonal cut at } x = a\}.$$

Informally, we may think of the line $x = a$ as determining the left- and right-hand sides of a multipartition in the obvious way. The catch is that these pieces won't be multipartitions, and so we add in as few nodes as possible to make them multipartitions.

Proposition 2.9. *Given $\lambda \in \mathcal{P}_n^\ell$, we have a bijection*

$$\begin{aligned} \Lambda_a(n, \theta, \kappa) &\xrightarrow{\sim} \Lambda_a^L(n_L, \theta, \kappa) \times \Lambda_a^R(n_R, \theta, \kappa) \\ \mu &\longmapsto (\mu^L, \mu^R). \end{aligned}$$

Proof. This is clear from the definitions. $\square$

Remark 2.10. Given $\lambda \in \mathcal{P}_n^\ell$ and $\mu \in \Lambda_a(n, \theta, \kappa)$, one can write down the ℓ -multipartitions μ^L and μ^R explicitly as follows. Suppose $\mu^{(m)} = (\mu_1, \mu_2, \dots)$. If every node in $[\mu^{(m)}]$ lies to the left of $a - g$ (respectively right of a), then the m th component of μ^L (respectively μ^R) is just $\mu^{(m)}$ itself.

Otherwise, at least one node in $[\mu^{(m)}]$ has x -coordinate in $(a - g, a)$. Suppose the highest such node is $(r, c, m) \in [\mu^{(m)}]$. Then the m th components of μ^L and μ^R are

$$(\mu^L)^{(m)} = (\mu_1, \mu_2, \dots, \mu_r) \quad \text{and} \quad (\mu^R)^{(m)} = (c^r, \mu_{r+1}, \mu_{r+2}, \dots).$$

Example 2.11. Continuing Example 2.2, recall that $e = 3$, $g = 2$, $\theta = 0$ and $(\lambda, \mu) = ((5, 4, 3, 2, 1), (4^3, 1^3))$. Making a diagonal cut at $x = 1$ yields

$$\lambda^L = (5, 4, 3), \quad \lambda^R = (3^3, 2, 1), \quad \mu^L = (4^3), \quad \mu^R = (3^3, 1^3).$$

These are depicted in Figure 5. We note that the only difference between our diagonal cuts and the more classical horizontal and vertical cuts is that we have an ‘extra rectangle’ of boxes in the form of a partition (r^c) common to *both* λ^L and λ^R .




FIGURE 5. Let (λ, μ) be the pair of partitions admitting a diagonal cut as in Figure 4. We picture the resulting partitions above. The leftmost diagram depicts λ^L, μ^L , and the rightmost diagram depicts λ^R, μ^R .

2.3. A subquotient of the diagrammatic Cherednik algebra. We now define the subquotients of the diagrammatic Cherednik algebras in which we shall be interested for the remainder of the paper.

Definition 2.12. Given $\lambda \in \mathcal{P}_n^\ell$ and $a \in \mathbb{R}$, we define idempotents

$$e = \sum_{\mu \in E \cap F} 1_\mu \quad \text{and} \quad f = \sum_{\mu \in F \setminus E} 1_\mu$$

in $A(n, \theta, \kappa)$. We let $A_{\Lambda_a(n, \theta, \kappa)}$ denote the subquotient of $A := A(n, \theta, \kappa)$ given by

$$e(A/(AfA))e.$$

Proposition 2.13. *The algebra $A_{\Lambda_a(n, \theta, \kappa)}$ is a graded cellular algebra with a highest weight theory. The cellular basis is given by*

$$\{C_{ST} \mid S \in \text{SStd}(\alpha, \beta), T \in \text{SStd}(\alpha, \gamma), \alpha, \beta, \gamma \in \Lambda_a(n, \theta, \kappa)\},$$

with respect to the θ -dominance order on $\Lambda_a(n, \theta, \kappa)$. The graded decomposition numbers and graded higher extension groups are preserved under taking this subquotient.

Proof. The set $F \setminus E$ is cosaturated in the θ -dominance ordering and so

$$\langle C_{ST} \mid S \in \text{SStd}(\lambda, \mu), T \in \text{SStd}(\lambda, \nu), \lambda \in F \setminus E, \mu, \nu \in \mathcal{P}_n^\ell \rangle_{\mathbb{C}}$$

is an ideal of $A(n, \theta, \kappa)$. The resulting quotient algebra has basis indexed by $S \in \text{SStd}(\alpha, \beta)$, $T \in \text{SStd}(\alpha, \gamma)$ such that $\alpha, \beta, \gamma \notin (F \setminus E)$ (by conditions (2) and (3) of Definition 1.1 and Theorem 1.12). Applying the idempotent truncation to this basis (and using (6) of Definition 1.1) we obtain the required basis of $A_{\Lambda_a(n, \theta, \kappa)}$. The graded decomposition numbers (as well as dimensions of higher extension groups) are preserved under both the quotient and truncation maps; this follows by the arguments of [Don98, Appendix: Lemmas A3.1 and A3.3] (for the ungraded case) as the ideal is generated by a (degree zero) idempotent. $\square$

Definition 2.14. Let $\mu, \nu \in \Lambda_a(n, \theta, \kappa)$. For $S \in \text{SStd}(\mu, \nu)$, we define S^L and S^R to be the tableaux determined by

- (1) $S^L(r, c, k) = S(r, c, k)$ for $\mathbf{i}_{(r,c,k)} < a$ (respectively $S^R(r, c, k) = S(r, c, k)$ for $\mathbf{i}_{(r,c,k)} > a$);
(2) $S^L(r, c, k) = \mathbf{i}_{(r,c,k)}$ for $\mathbf{i}_{(r,c,k)} > a$ (respectively $S^R(r, c, k) = \mathbf{i}_{(r,c,k)}$ for $\mathbf{i}_{(r,c,k)} < a$);
for $(r, c, k) \in [\mu^L]$ (respectively $(r, c, k) \in [\mu^R]$).

Theorem 2.15. *Given $\lambda \in \mathcal{P}_n^\ell$ and $a \in \mathbb{R}$. We have an isomorphism*

$$A_{\Lambda_a(n, \theta, \kappa)} \cong A_{\Lambda_a^L(n_L, \theta, \kappa)} \otimes_{\mathbb{k}} A_{\Lambda_a^R(n_R, \theta, \kappa)}$$

of graded vector spaces over $\mathbb{k}$ given by the map

$$\varphi : C_{\mathbf{ST}} \mapsto C_{S^L T^L} \otimes C_{S^R T^R}.$$

Proof. Recall from Proposition 2.9 that the map $\Lambda_a(n, \theta, \kappa) \cong \Lambda_a^L(n_L, \theta, \kappa) \times \Lambda_a^R(n_R, \theta, \kappa)$ given by $\mu \mapsto \mu^L \times \mu^R$, is bijective. Given $\mu, \nu \in \Lambda_a(n, \theta, \kappa)$, we have that

$$\begin{aligned} |D_\mu \cap (-\infty, a - g)| &= |D_\nu \cap (-\infty, a - g)|, \\ |D_\mu \cap (a - g, a)| &= |D_\nu \cap (a - g, a)|, \\ |D_\mu \cap (a, \infty)| &= |D_\nu \cap (a, \infty)| \end{aligned}$$

and therefore (by Definition 1.6) any tableau $S \in \text{SStd}(\mu, \nu)$ satisfies $S(r, c, k) = (r', c', k')$ for $\mathbf{i}_{(r,c,k)}$ and $\mathbf{i}_{(r',c',k')}$ a pair of points in the same region: $(-\infty, a - g)$, $(a - g, a)$, or (a, ∞) . Therefore $S^L \in \text{SStd}(\mu^L, \nu^L)$, $S^R \in \text{SStd}(\mu^R, \nu^R)$, and the map $S \mapsto S^L \times S^R$ is a bijection $\text{SStd}(\mu, \nu) \rightarrow \text{SStd}(\mu^L, \nu^L) \times \text{SStd}(\mu^R, \nu^R)$. Moreover, we have that

$$(2.4) \quad \begin{aligned} B_{S^L} \cap (-\infty, a - g) \times [0, 1] &= B_S \cap (-\infty, a - g) \times [0, 1] \\ B_{S^R} \cap (a, \infty) \times [0, 1] &= B_S \cap (a, \infty) \times [0, 1]. \end{aligned}$$

Given any two multipartitions $\mu^L, \nu^L \in \Lambda_a^L(n_L, \theta, \kappa)$ (respectively $\mu^R, \nu^R \in \Lambda_a^R(n_R, \theta, \kappa)$) all nodes to the right of the point $a - g$ (respectively to the left of a) are common to both multipartitions. Therefore by the definition of semistandard tableau, we have that $T(r, c, k) = \mathbf{i}_{(r,c,k)}$ for $\mathbf{i}_{(r,c,k)} > a - g$ (respectively $T(r, c, k) = \mathbf{i}_{(r,c,k)}$ for $\mathbf{i}_{(r,c,k)} < a$) for all $T \in \text{SStd}(\mu^L, \nu^L)$ (respectively, for all $T \in \text{SStd}(\mu^R, \nu^R)$). Therefore, we have that

$$(2.5) \quad \begin{aligned} B_{S^L} \cap ((a - g, \infty) \times [0, 1]) &= 1_\mu \cap ((a - g, \infty) \times [0, 1]) = 1_\nu \cap ((a - g, \infty) \times [0, 1]) \\ B_{S^R} \cap ((-\infty, a) \times [0, 1]) &= 1_\mu \cap ((-\infty, a) \times [0, 1]) = 1_\nu \cap ((-\infty, a) \times [0, 1]) \end{aligned}$$

and the diagrams in equation (2.5) are both of degree zero. Therefore by equations (2.4) and (2.5) we have that

$$\deg(B_{S^L}) + \deg(B_{S^R}) = \deg(B_S \cap (-\infty, a - g) \times [0, 1]) + \deg(B_S \cap (a, \infty) \times [0, 1]) = \deg(B_S)$$

and the map $B_S \mapsto B_{S^L} \times B_{S^R}$ is a degree preserving bijection on the cell modules. The result follows. $\square$

Example 2.16. Recall our running example, with $e = 3$, $g = 2$, $\theta = 0$ and $(\lambda, \mu) = ((5, 4, 3, 2, 1), (4^3, 1^3))$. There is a unique element $S \in \text{SStd}(\lambda, \mu)$ and the corresponding basis element B_S is depicted in Figure 6. This pair admits a θ -diagonal cut at $x = 1$. The elements B_{S^L} and B_{S^R} are depicted in Figure 7.

Theorem 2.17. *The map $\varphi : A_{\Lambda_a(n, \theta, \kappa)} \rightarrow A_{\Lambda_a^L(n_L, \theta, \kappa)} \otimes_{\mathbb{k}} A_{\Lambda_a^R(n_R, \theta, \kappa)}$, defined in Theorem 2.15, is an isomorphism of graded $\mathbb{k}$ -algebras.*

Proof. We have already seen in Theorem 2.15 that the map φ is an isomorphism of graded vector spaces over $\mathbb{k}$; it remains to check that $C_{S^L T^L} C_{U^L V^L} \otimes C_{S^R T^R} C_{U^R V^R} = \varphi(C_{S T}) \varphi(C_{U V}) = \varphi(C_{S T} C_{U V})$. The cell-ideals of $A(n, \theta, \kappa)$ are ordered according to the θ -dominance order on loadings; more dominant loadings in this ordering are given by moving strands to the left. Therefore in order to rewrite a product in terms of the cellular basis, we shall proceed by pulling the strands in the diagrams as far to the left as possible using relations (2.1) to (2.15).

The relations of $A(n, \theta, \kappa)$ (and therefore those of the subquotients $A_{\Lambda_a(n, \theta, \kappa)}$, $A_{\Lambda_a^L(n_L, \theta, \kappa)}$, and $A_{\Lambda_a^R(n_R, \theta, \kappa)}$) can all be applied to the regions $(-\infty, a - g)$, then $(a - g, a)$, and then (a, ∞) to each of the diagrams in turn. This is because relations (2.1) to (2.14) of $A(m, \theta, \kappa)$ for $m = n, n_L, n_R$




FIGURE 6. The basis element B_S corresponding to $(\mu, \nu) = ((5, 4, 3, 2, 1), (4^3, 1^3))$. Given $a = 1$, we mark the lines of the cut, $x = a$ and $x = a - g$.




FIGURE 7. The basis elements B_{S^L} and B_{S^R} corresponding to $(\mu, \nu) = ((5, 4, 3, 2, 1), (4^3, 1^3))$ cut at $x = 1$ and the unique $S \in \text{SStd}(\mu, \nu)$.

are local, and the additional idempotent relations (relation (2.15) and the cosaturated quotient relation in the θ -dominance order) are all given in terms of loadings, and therefore are compatible with restriction to regions of $\mathbb{R}$.

First, apply relations (2.1) to (2.15) to the region $(-\infty, a - g)$ of the diagrams $C_{ST}C_{UV}$ and $C_{S^L T^L}C_{U^L V^L}$ concurrently to push all strands in $C_{ST}C_{UV} \cap ((-\infty, a - g) \times [0, 1])$ and $C_{S^L T^L}C_{U^L V^L} \cap ((-\infty, a - g) \times [0, 1])$ as far to the left as possible.

Next, apply relations (2.1) to (2.15) to the region (a, ∞) of the diagrams $C_{ST}C_{UV}$ and $C_{S^R T^R}C_{U^R V^R}$ concurrently to push all strands in $C_{ST}C_{UV} \cap ((a, \infty) \times [0, 1])$ and $C_{S^R T^R}C_{U^R V^R} \cap ((a, \infty) \times [0, 1])$ as far to the left as possible.

If every element in the resulting linear combination of diagrams in $A(n, \theta, \kappa)$ is an element of the basis $\mathcal{C}$, then the result follows. Otherwise, let d be any diagram in the linear combination which is not in $\mathcal{C}$. By assumption, the diagram d contains a strand which cannot be pushed further left (using relations (2.1) to (2.15)) in either $(a - g, a)$ or (a, ∞) , but which can be pushed further left in $(-\infty, \infty)$. In the former (respectively latter) case, pull this strand to the left of $a - g$ (respectively a); the resulting diagram factors through an idempotent with loading $\mathbf{i}_\nu$, for some ν such that $|L_a(\nu)| > |L_a(\lambda)|$ (respectively $|I_a(\nu)| + |L_a(\nu)| > |I_a(\lambda)| + |L_a(\lambda)|$), and so is zero in $A_{\Lambda_a(n, \theta, \kappa)}$. (Here we have implicitly made use of Lemma 1.11.)

Now consider the element $\varphi(d) = d' \otimes d'' \in A(n_L, \theta, \kappa) \otimes A(n_R, \theta, \kappa)$. In the former (respectively latter) case, there is a strand in the diagram $d' \in A(n_L, \theta, \kappa)$ (respectively $d'' \in A(n_R, \theta, \kappa)$) which can be pulled to the left of $a - g$ (respectively a); the resulting diagram factors through an idempotent with loading $\mathbf{i}_\nu$, for some $\nu \in \mathcal{P}_{n_L}^\ell$ (respectively for some $\nu \in \mathcal{P}_{n_R}^\ell$) such that $|L_a(\nu)| > |L_a(\lambda^L)|$ (respectively $|I_a(\nu)| + |L_a(\nu)| > |I_a(\lambda^R)| + |L_a(\lambda^R)|$) and so is zero

in $A_{\Lambda_a^L(n_L, \theta, \kappa)}$ (respectively $A_{\Lambda_a^R(n_R, \theta, \kappa)}$). (Here we have implicitly made use of Lemma 1.11.) The result follows. $\square$

Corollary 2.18. *Let (λ, μ) be a pair of ℓ -multipartitions of n such that $\mu \leq_\theta \lambda$. If, for some $a \in \mathbb{R}$, (λ, μ) admits a θ -diagonal cut at $x = a$, then*

$$d_{\lambda\mu}(t) = d_{\lambda^L \mu^L}(t) \times d_{\lambda^R \mu^R}(t)$$

and, furthermore,

$$\text{Ext}_{A(n, \theta, \kappa)}^k(\Delta(\lambda), \Delta(\mu)) \cong \bigoplus_{i+j=k} \text{Ext}_{A(n_L, \theta, \kappa)}^i(\Delta(\lambda^L), \Delta(\mu^L)) \otimes \text{Ext}_{A(n_R, \theta, \kappa)}^j(\Delta(\lambda^R), \Delta(\mu^R)),$$

where $n_L = |\lambda^L| = |\mu^L|$ and $n_R = |\lambda^R| = |\mu^R|$.

Proof. This follows by Proposition 2.13 and Theorem 2.17. $\square$

Remark 2.19. As previously noted, in the level 1 case, a pair (λ, μ) such that $\lambda \geq \mu$ admits a θ -diagonal cut at $x = a$ if and only if it admits a horizontal cut, if and only if it admits a vertical cut. These horizontal and vertical cuts cross at the ‘highest’ node in the diagonal $D_\lambda \cap (a - g, a)$; that is, the node (r, c) satisfying equation (2.1) for which $r + c$ is maximal. In this case, the corresponding horizontal and vertical cuts are after the r th row and the c th column, respectively. In this case, the ungraded version of the result above is a corollary of [Don98, 4.2(9)] and [Don07, Section 10]. (We remark that the graded case is new, although the result for graded decomposition numbers was already known over $\mathbb{C}$ by calculations in the Fock-space – see [CMT02]).

For (graded) decomposition numbers and higher extension groups of cyclotomic q -Schur algebras (in other words, the algebras $A(n, \theta, \kappa)$ for θ a well-separated weighting) the result above is completely new. In this case it is easy to see that a pair of multipartitions (λ, μ) admits a θ -diagonal cut at $x = a$ if and only if it admits a horizontal cut after the r th row of the m th component (in the sense of [FS]), where the node (r, c, m) is the highest node in $(a - g, a)$, as in Remark 2.10. Thus we retrieve an analogue of [FS, Theorem 4.8] (which is stated for the cyclotomic Hecke algebras) for the cyclotomic q -Schur algebras. Here, our ‘left piece’ is the same as the ‘top piece’ in [FS, Theorem 4.8] (modulo empty components which do not play an important role where these decomposition numbers are concerned), while our right piece differs from their ‘bottom piece’ in component m ; we have an extra r rows of length c , which are common λ^R and μ^R . In order to remove this ‘extra rectangle’ it would suffice to establish a ‘first row removal’ isomorphism of algebras relating algebras of different degrees and *more importantly* different e -multicharges and weightings; this would proceed by analogy with [FS]. Given an algebra with a well-separated weighting, it is not difficult to establish such an isomorphism on the level of graded vector spaces and thus deduce the equality of graded decomposition numbers over fields of characteristic zero (by arguing in an LLT-type fashion as in [BS, Theorem 4.12]). However, the obvious map does not lift to the level of an isomorphism of algebras and so we do not include this here.

Example 2.20. Recall our running example, with $e = 3$, $g = 2$, $\theta = 0$, and $(\lambda, \mu) = ((5, 4, 3, 2, 1), (4^3, 1^3))$. We have that

$$d_{(5,4,3,2,1)(4^3,1^3)}(t) = d_{(5,4,3)(4^3)}(t) d_{(3^3,2,1)(3^3,1^3)}(t) = t \times t = t^2.$$

regardless of the characteristic of the underlying field $\mathbb{k}$. Similarly,

$$\begin{aligned} & \text{Dim}_t(\text{Hom}_{A(15, (0), (0))}(\Delta(5, 4, 3, 2, 1), \Delta(4^3, 1^3))) \\ &= \text{Dim}_t(\text{Hom}_{A(12, (0), (0))}(\Delta(5, 4, 3), \Delta(4^3))) \times \text{Dim}_t(\text{Hom}_{A(12, (0), (0))}(\Delta(3^3, 2, 1), \Delta(3^3, 1^3))) \\ &= t \times t = t^2. \end{aligned}$$

Example 2.21. Let $e = 5$, $\kappa = (0, 2)$, $g = 1$, and $\theta = (0, 0.5)$ a FLOTW weighting. The bipartitions

$$\lambda = ((11, 9, 7, 3^2, 2, 1^3), (9, 4, 2, 1^4)) \quad \text{and} \quad \mu = ((10, 9, 8, 4, 3, 1^5), (8, 4, 2, 1^4))$$

admit a diagonal cut at $x = 2.6$, which yields

$$\begin{aligned}\lambda^L &= ((11, 9, 7, 3^2), (9, 4, 2)), & \lambda^R &= ((3^5, 2, 1^3), (1^7)), \\ \mu^L &= ((10, 9, 8, 4, 3), (8, 4, 2)), & \mu^R &= ((3^5, 1^5), (1^7)).\end{aligned}$$

These bipartitions are depicted in Figures 1 and 2 in the introduction. Applying Corollary 2.18, we have

$$d_{\lambda\mu}(t) = d_{\lambda^L\mu^L}(t) \times d_{\lambda^R\mu^R}(t).$$

First of all, we note that [BS, Theorem 4.30] is insufficient to calculate $d_{\lambda\mu}(t)$; this is because λ and μ differ in nodes of adjacent residues (0 and 1). However, [BS, Theorem 3.8] is sufficient to calculate both $d_{\lambda^L\mu^L}(t)$ and $d_{\lambda^R\mu^R}(t)$, because λ^L and μ^L (respectively λ^R and μ^R) differ only in nodes of a single residue, namely 0 (respectively 1). Applying [BS, Theorem 3.8], we deduce that

$$d_{\lambda\mu}(t) = (t^5 + t^3) \times t^2 = t^7 + t^5.$$

Example 2.22. Let $e = 3$, $\kappa = (0, 1)$, $g = 2$ and $\theta = (0, 115)$ a well-separated weighting. The bipartitions $\lambda = ((5^2, 4^2, 3, 2, 1), (9, 6, 4^2, 3, 2^3, 1))$ and $\mu = ((5, 4^2, 3^3), (9, 6, 5, 4^2, 2^2, 1^3))$ of 57 admit a θ -diagonal cut at $x = 121$. This cut results in

$$\begin{aligned}\lambda^L &= ((5^2, 4^2, 3, 2, 1), (9, 6, 4^2, 3)), & \lambda^R &= (\emptyset, (3^5, 2^3, 1)), \\ \mu^L &= ((5, 4^2, 3^3), (9, 6, 5, 4^2)), & \mu^R &= (\emptyset, (3^5, 2^2, 1^3)).\end{aligned}$$

This reduction again yields multipartitions amenable to the techniques of [BS] (whereas λ and μ are not). The left-hand pieces should be compared with [BS, Example 2.6], which yields $d_{\lambda^L\mu^L} = t^{11} + 2t^9 + 2t^7 + t^5$. We may also apply [BS, Theorem 4.30] to calculate that $d_{\lambda^R\mu^R} = t$. Thus, we have $d_{\lambda\mu} = t^{12} + 2t^{10} + 2t^8 + t^6$.

REFERENCES

- [BCS] C. Bowman, A. Cox, and L. Speyer, *On graded decomposition numbers for rational Cherednik algebras*, [arXiv:1503.07088](#), 2015, preprint. [Page 7.]
- [BS] C. Bowman and L. Speyer, *Kleshchev’s decomposition numbers for diagrammatic Cherednik algebras*, [arXiv:1507.06631](#), 2015, preprint. [Pages 1, 2, 14, 15.]
- [CMT02] J. Chuang, H. Miyachi, and K. M. Tan, *Row and column removal in the q -deformed Fock space*, J. Algebra **254** (2002), no. 1, 84–91. [Pages 1, 14.]
- [DJM98] R. Dipper, G. D. James, and A. Mathas, *Cyclotomic q -Schur algebras*, Math. Z. **229** (1998), no. 3, 385–416. [Page 1.]
- [Don85] S. Donkin, *A note on decomposition numbers for general linear groups and symmetric groups*, Math. Proc. Cambridge Philos. Soc. **97** (1985), no. 1, 57–62. [Pages 1, 9.]
- [Don98] S. Donkin, *The q -Schur algebra*, London Mathematical Society Lecture Note Series, vol. 253, Cambridge University Press, Cambridge, 1998. [Pages 11, 14.]
- [Don07] ———, *Tilting modules for algebraic groups and finite dimensional algebras*, Handbook of tilting theory, London Math. Soc. Lecture Note Ser., vol. 332, Cambridge Univ. Press, Cambridge, 2007, pp. 215–257. [Pages 1, 14.]
- [FL03] M. Fayers and S. Lyle, *Row and column removal theorems for homomorphisms between Specht modules*, J. Pure Appl. Algebra **185** (2003), 147–164. [Pages 1, 9.]
- [FS] M. Fayers and L. Speyer, *Generalised column removal for graded homomorphisms between Specht modules*, [arXiv:1404.4415](#), 2014, preprint. [Page 14.]
- [HM10] J. Hu and A. Mathas, *Graded cellular bases for the cyclotomic Khovanov–Lauda–Rouquier algebras of type A*, Adv. Math. **225** (2010), 598–642. [Page 3.]
- [Jam81] G. D. James, *On the decomposition matrices of the symmetric groups, III*, J. Algebra **71** (1981), no. 1, 115–122. [Page 1.]
- [LM05] S. Lyle and A. Mathas, *Row and column removal theorems for homomorphisms of Specht modules and Weyl modules*, J. Algebraic Combin. **22** (2005), 151–179. [Page 1.]
- [Los] I. Losev, *Proof of Varagnolo–Vasserot conjecture on cyclotomic categories $\mathcal{O}$* , Selecta Math. (to appear). [Page 1.]
- [RSVV] R. Rouquier, P. Shan, M. Varagnolo, and E. Vasserot, *Categorifications and cyclotomic rational double affine Hecke algebras*, Invent. Math. (to appear). [Page 1.]
- [Web] ———, *Knot invariants and higher representation theory*, Mem. Amer. Math. Soc. (to appear). [Page 1.]

- [Webb] B. Webster, *Weighted Khovanov–Lauda–Rouquier algebras*, [arXiv:1209.2463v3](#), 2015, preprint. [Page 8.]
- [Webc] B. Webster, *Rouquier’s conjecture and diagrammatic algebra*, [arXiv:1306.0074v3](#), 2014, preprint. [Pages 1, 5, 6, 8.]

E-mail address: `Chris.Bowman.2@city.ac.uk`

DEPARTMENT OF MATHEMATICS, CITY UNIVERSITY LONDON, NORTHAMPTON SQUARE, LONDON, EC1V 0HB, UK

E-mail address: `l.speyer@ist.osaka-u.ac.jp`

OSAKA UNIVERSITY, SUITA, OSAKA, 565-0871, JAPAN